



Recommendations for general surgery activities in a pandemic scenario (SARS-CoV-2)

F. Di Marzo¹, M. Sartelli², R. Cennamo¹, G. Toccafondi³, F. Coccolini⁴, G. La Torre⁵, G. Tulli⁶, M. Lombardi⁷ and M. Cardì⁸

¹UOC Chirurgia Generale, Ospedale Valtiberina – Sansepolcro, Usl Toscana Sud-Est, ²UOC Chirurgia Generale, Ospedale di Macerata, ³Centro Gestione Rischio Clinico, Toscana, ⁴UOC Chirurgia Generale e d'urgenza, AOU Pisa, ⁵Dipartimento di Sanità Pubblica e Malattie Infettive, “Sapienza” Università di Roma, ⁶Agenzia Regionale di Sanità Toscana, Firenze, ⁷UOC Chirurgia Generale, NOA – Marina di Massa, Usl Toscana Nord-Ovest, ⁸Dipartimento di Chirurgia “P. Valdoni”, “Sapienza” Università di Roma, Italy

Correspondence to: Dr F. Di Marzo, UOC Chirurgia Generale, Ospedale Valtiberina – Sansepolcro, Usl Toscana Sud-Est – Viale Galileo Galilei, 101 – 52037 Sansepolcro (AR), Italy (e-mail: fdimarzo@yahoo.com)

Paper accepted 3 April 2020

Published online in Wiley Online Library (www.bjs.co.uk). DOI: 10.1002/bjs.11652

The SARS-CoV-2 pandemic has over 1 million cases with more than 50 000 deaths globally to date (early April 2020)¹. Disease spread in Italy has been rapid, with in excess of 115 000 SARS-CoV-2-positive patients and 13 000 deaths (1 April 2020). Mortality is currently 5.2 per cent in Tuscany, compared with 11 per cent nationwide. Healthcare systems have been forced to reorganize surgical activities² as they struggle with an overwhelming number of critically ill patients. One of the lessons we learnt from the 2003 SARS outbreak is that case identification was limited by non-specific mild symptoms, with a flu-like prodrome progressing to fulminant disease, which could quickly overwhelm hospitals³ without plans for surge capacity, patient transfer, or sufficient numbers of trained staff. In addition to this, initial delays in transposing regional indications to stop surgical activities and to implement proper infection control procedures contributed to a disproportionate number of healthcare workers becoming infected (approximately 9000 in Italy alone by 30 March 2020).

The organizational model adopted was based on a rapid response to early signs of incoming crisis, and elective surgical activity (including outpatient consultations) was reduced to a minimum. Hospital areas capable of intensive care unit (ICU) level monitoring, including the post anaesthesia care unit or operating room, were repurposed as temporary ICU facilities. On 6 March the regional government defined three types of hospitals. Type A university hospitals are focused mainly on SARS-CoV-2 patient care, whereas type B hub-hospitals have ICU resources and the ability to organize two different pathways for SARS-CoV-2-positive and -negative patients; in both type A and B hospitals, scheduled elective surgery and outpatient activities are suspended, while emergency and urgent

surgery, cancer treatments, transplants and burn care centres are running normally. Any admitted patient must be tested for SARS-CoV-2. Type C spoke hospitals with limited or no ICU beds, including private hospitals, are dedicated to non-SARS-CoV-2 patients.

Infected patients admitted via the emergency room should be quarantined (both spatially and in staff resources) according to the local infectious diseases policy. To decrease the risk of in-hospital transmission, patients with SARS-CoV-2 should be separated from others, ideally in single rooms, and staff should consider wearing surgical or N95 masks, a disposable cap to cover the hair, and gloves⁴. Medical personnel should use fast-drying hand antiseptics and change gloves immediately after contact with a patient, body fluids or contaminated material⁵. Elective surgical activity is reduced to a minimum so operating rooms can be converted into ICUs using post anaesthesia care unit beds first, leaving space for emergency/urgent surgery. Admissions should follow two different pathways: one for SARS-CoV-2-positive and the other for -negative patients. To decrease the risk of in-hospital transmission hospitals should separate staff caring for patients with SARS-CoV-2 from others. All SARS-CoV-2-negative patients should also have the shortest possible hospital stay and be cared for in a designated ‘infection-free’ area or spoke (non-SARS-CoV-2) hospital.

High-intensity, pressured work will lead to staff fatigue and even psychological crises so staff need psychological support, reasonable scheduling, and appropriate rest. First-line medical staff should try to identify their negative psychological emotions, enable positive self-counselling, and seek professional help when needed⁶. Symptoms and/or fever should stimulate testing for the virus expeditiously. Ideally, there should be routine testing of healthcare

staff, serological confirmation immunity, and vaccination of those without natural immune responses.

The numbers of emergency admissions with coexistent viral infection are increasing, adding to the pressure on resources. Experienced surgeons should assess patients before admission to reduce unnecessary admissions by 20 to 30 per cent, with substantial resource savings⁷. Effective communication should ensure closed-loop information transfer on any patient's SARS-CoV-2 status. Transanal procedures, stomas for faecal diversion, and intestinal anastomoses may risk aerosol contamination⁸. Laparoscopy is an option but the relative risk of viral exposure *versus* open procedures remains unclear and in critically ill patients with lung dysfunction, sepsis or shock, open surgery is advised. An operating room maintained at negative pressure with a high frequency air exchanges (at least 25 cycles/h) or designated 'contaminated' surgery-only is sensible⁹.

Surgical patients who are suspected of being or who test positive for SARS-CoV-2 infections should follow the local management protocol, which may include: wearing specified bracelets and surgical masks, having their medical records marked with warning labels, and being brought through defined routes and lifts to the special isolation area for recovery. A dedicated portable ventilator should be used when managing infected ICU patients, switching off the gas flow and closing the endotracheal tube to reduce aerosol production when connecting to the operating room ventilator¹⁰. The gas sampling tube should be protected by a high-efficiency particulate air (HEPA) filter and the soda lime should be changed regularly¹⁰. Anaesthetists should use three-level protective measures with HEPA filters on both limbs of the circuit and consider disposable components⁹. For surgical cases in infected patients, all staff should take a training course on personal protective equipment use: disposable double-layer hats, N95 masks, surgical gowns for single use or medical protective clothing, goggles, or full face mask, and double-layer sterile gloves. According to the Association for the Advancement of Medical Instrumentation (AAMI), the protection level of the surgical gowns depends on the type of procedure. Level 4 is the highest fluid and microbial barrier, and is needed for long, fluid-intensive procedures; level 3 is indicated as a moderate fluid barrier protection¹¹.

The Chinese New Coronavirus Infected Pneumonia Diagnosis and Treatment Plan (Trial Version 7) highlighted the following criteria for considering removing quarantine and transferring the patient to a general ward¹²: a normal body temperature for more than 3 days; respiratory symptoms significantly relieved; lung imaging showing that the inflammation has been significantly absorbed; and a negative SARS-CoV-2 nucleic acid

amplification test (sampling interval at least 24 h) for specialist treatment or discharge based on their condition. The Italian Ministero della salute – Consiglio Superiore di Sanità stated on 28 February 2020 that SARS-CoV-2 patients can be considered cured if the symptoms resolve and two tests for SARS-CoV-2 at 24-h intervals are negative. For patients who clinically recover earlier than 7 days after onset, an interval of 7 days should elapse between the first and the final test. Virus clearance is defined as viral RNA disappearance from bodily fluids accompanied by the appearance of specific IgG.

The Singapore National Centre for Infectious Diseases (NCID) released the following guidelines on de isolation of SARS-CoV-2 suspected cases¹³: discharge patient with advisory and clinical follow-up if indicated and with daily calls until day 14 after the last possible exposure, under the following conditions: patient afebrile for at least 24 h, two respiratory samples tested negative for SARS-CoV-2 by polymerase chain reaction (PCR) in at least 24 h, day of illness from onset at least 6 days, or alternative aetiology found (for example, influenza, bacteraemia), or no close contact with a SARS-CoV-2 case, or no in-patient care required for other reasons. The US Centre for Disease Control released interim guidance for discontinuation of transmission-based precautions and disposition of hospitalized patients with SARS-CoV-2¹⁴: negative real-time PCR results from at least two consecutive sets of nasopharyngeal and throat swabs collected at least 24 h apart from a patient with SARS-CoV-2 (a total of four negative specimens) and resolution of fever, without antipyretic medications, improvement in illness signs and symptoms.

Regarding possible antiviral therapy, the Italian Society for Infectious and Tropical Disease released guidelines on 13 March 2020. The cut-off is the condition of the respiratory system evaluated using the Brescia Covid Respiratory severity scale: in patients before intubation the disease is primarily controlled by anti-inflammatory treatment (tocilizumab) in association with early antiviral and antiretroviral therapy. In mechanically ventilated patients the strategy is based on high-pressure ventilation and prone positioning. There is no evidence of efficacy of prophylactic chloroquine or hydroxychloroquine and the use is not recommended. Remdesvir has been shown to be active in preclinical studies on SARS-CoV and MERS-CoV infections by acting on the viral polymerase of coronavirus. The drug is not registered in Italy, and can be used only for 'compassionate' reasons, subject to approval by an Ethics Committee. There are multiple clinical trials ongoing and it is expected that a vaccine will be available within months.

Acknowledgements

The authors thank Alice Crossman for professional editing and language revision.

Disclosure: The authors declare no conflict of interest.

References

- 1 World Health Organization. *Coronavirus disease (COVID-19) outbreak webpage*. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
- 2 Spinelli A, Pellino G. COVID-19 pandemic: perspectives on an unfolding crisis. *Br J Surg* 2020; doi: 10.1002/bjs.11627 [Epub ahead of print].
- 3 Cheng VC, Chan JF, To KK Yuen KY. Clinical management and infection control of SARS: lessons learned. *Antiviral Res* 2013; **100**: 407–419.
- 4 Family Health Committee of the People's Republic of China New. *Coronavirus Transmission Routes and Prevention Guidelines* [EB/OL], (2020-01-27) [2020-02-12].
- 5 Family Health Committee of the People's Republic of China. *Hand Hygiene Rules for Medical Staff* [EB/OL]. (2009-04-01) [2020-02-12].
- 6 Family Health Committee of the People's Republic of China. *About the Pneumonia Epidemic of New Coronavirus Infection. Notice of Guiding Principles for Intervention of Emergent Psychological Crisis* [EB/OL] (2020-01-26) [2020-02-12].
- 7 Anderson I. *The Future of Emergency General Surgery - A Joint Document*; 2015. https://www.acpgbi.org.uk/content/uploads/2016/07/Future-of-EGS-joint-document_Iain-Anderson_140915.pdf.
- 8 Yu GY, Lou Z, Zhang W. Several suggestions of operation for colorectal cancer under the outbreak of Corona Virus Disease 19 in China. *Zhonghua Wei Chang Wai Ke Za Zhi* 2020; **23**: 9–11. doi:10.3760/cma.j.issn.1671-0274.2020.03.002 [E-pub ahead of print].
- 9 Wong J, Goh QY, Tan Z, Lie SA, Tay YC, Ng SY, Soh CR. Preparing for a COVID-19 pandemic: a review of operating room outbreak response measures in a large tertiary hospital in Singapore. *Can J Anaesth*. 2020; doi: 10.1007/s12630-020-01620-9 [Epub ahead of print].
- 10 Ti LK, Ang LS, Foong TW, Ng BSW. What we do when a COVID-19 patient needs an operation: operating room preparation and guidance. *Can J Anaesth*. 2020; doi: 10.1007/s12630-020-01617-4 [Epub ahead of print].
- 11 Association for the Advancement of Medical Instrumentation. *Liquid Barrier Performance and Classification of Protective Apparel and Drapes Intended for Use in Health Care Facilities*; 2012 (ANSI/AAMI PB70:2012):6–7 (§4.2.1–4.2.3).
- 12 People's Republic of China National Health Council State Administration of Traditional Chinese Medicine. *New Coronavirus Infection Pneumonia Diagnosis and Treatment Plan* [EB/OL] (2020-02-05) [2020-02-11].
- 13 Tay JY, Lim PL, Marimuthu K, Sadarangani SP, Ling LM, Ang BSP *et al.* e-isolating Coronavirus Disease 2019 Suspected Cases: A Continuing Challenge. *Clin Infec Dis* 2020, ciaa179, <https://doi.org/10.1093/cid/ciaa179>.
- 14 Center for Disease Control and Prevention. *Interim Guidance for Discontinuation of Transmission-Based Precautions and Disposition of Hospitalized Patients with COVID-19*; 2020. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html>.

The *BJS* team wish to reach out to express our support and gratitude to surgeons and healthcare workers around the globe. These are difficult times and your leadership is key to providing the best care possible. *BJS* welcomes submissions relating to the challenges faced in this pandemic (expect publication within a week). A blog has been launched (cuttingedgeblog.com) and publication of accepted pieces will be within hours.

Best wishes to you all.

DesWinter MD (Editor-in-Chief) on behalf of the *BJS* Editors, production team, Editorial Council and Board